

Eastern Kentucky University

Encompass

Psychology Faculty and Staff Research

Psychology

1-1-2012

Empirically Guided Case Conceptualization of Posttraumatic Stress Disorder with the Minnesota Multiphasic Personality Inventory-2 Restructured Form (MMPI-2-RF) in a Forensic Disability Evaluation

Lauren M. Vines
University of Louisville

Roger O. Gervais
The University of Alberta

Dustin Wygant
Eastern Kentucky University, dustin.wygant@eku.edu

Follow this and additional works at: https://encompass.eku.edu/psychology_fsresearch



Part of the [Clinical Psychology Commons](#)

Recommended Citation

Vines, Lauren M.; Gervais, Roger O.; and Wygant, Dustin, "Empirically Guided Case Conceptualization of Posttraumatic Stress Disorder with the Minnesota Multiphasic Personality Inventory-2 Restructured Form (MMPI-2-RF) in a Forensic Disability Evaluation" (2012). *Psychology Faculty and Staff Research*. 8. https://encompass.eku.edu/psychology_fsresearch/8

This Article is brought to you for free and open access by the Psychology at Encompass. It has been accepted for inclusion in Psychology Faculty and Staff Research by an authorized administrator of Encompass. For more information, please contact Linda.Sizemore@eku.edu.

Empirically Guided Case Conceptualization of Post-traumatic Stress Disorder with the Minnesota Multiphasic Personality Inventory-2 Restructured Form (MMPI-2-RF) In a Forensic Disability Evaluation

LAUREN M. VINES, M.S.
University of Louisville

DUSTIN B. WYGANT, PH.D.
Eastern Kentucky University

ROGER O. GERVAIS, PH.D.
Neurobehavioural Associates, Edmonton, Alberta
The University of Alberta

Abstract

The following article discusses how the Restructured Form of the Minnesota Multiphasic Personality Inventory (MMPI-2-RF; Ben-Porath & Tellegen, 2008) can be used in case conceptualizations for Posttraumatic Stress Disorder (PTSD), particularly in compensation seeking settings. We review contemporary conceptualizations of PTSD, particularly emphasizing the role that affect and personality in regards to etiology of the disorder, as well as different manifestations of the disorder. We then review the case of an

Author Notes

Lauren M. Vines, M.S., University of Louisville; Dustin B. Wygant, Ph.D., Department of Psychology, Eastern Kentucky University; Roger O. Gervais, Ph.D., Neurobehavioural Associates, Edmonton, Alberta. This case study was approved by the Institutional Review Board at Eastern Kentucky University. Please send any correspondence to Dr.

Dustin Wygant, Department of Psychology, Eastern Kentucky University. 521 Lancaster Ave. Richmond, KY 40475. Fax (859) 622-5871. Dustin.wygant@eku.edu.

(Abstract continued)

individual seeking compensation for trauma related disability performed by the third author. Particular emphasis is placed on examining how interpretation of the MMPI-2-RF profile is guided by empirical findings.

Empirically Guided Case Conceptualization of Posttraumatic Stress Disorder with the Minnesota Multiphasic Personality Inventory-2 Restructured Form (MMPI-2-RF) in Forensic Disability Evaluations

PTSD is a psychological condition that readily lends itself to presentation within the arena of forensic psychology (Resnick, West, & Payne, 2008). The very nature of its etiology requires an individual to experience or witness a potentially life threatening event (APA, 2000) that can result in physical and mental injuries for which monetary compensation may be pursued, whether they occur during military service or as a result of motor vehicle or other accidents. Consequently, PTSD symptoms are common during disability claims (Arbisi, Ben-Porath, & McNulty, 2006; Rubenzer, 2009). Experts have recommended a multi-method approach to the assessment of PTSD, emphasizing the assessment of malingering and genuine symptom report (see Resnick, West, & Payne, 2008; Rubenzer, 2009). Consequently, self-report measures such as the Minnesota Multiphasic Personality Inventory-2 (MMPI-2; Butcher, et al., 2001) have played a major role in the assessment of PTSD, particularly in forensic set-

tings, where clinicians must often establish the validity of a patient's symptom reports. A benefit of this instrument is its ability to identify several threats to protocol validity in the context of a forensic evaluation, such as random responding, acquiescent responding, as well as over and under-reporting of symptoms and problems, which has been extensively researched (e.g., Arbisi, Ben-Porath, & McNulty, 2006; Rogers et al., 2003). Previous studies have supported the use of these validity scales, namely the Infrequency (F) scale, Infrequency Back (Fb) scale, Infrequency Psychopathology (Fp) scale, and the Symptom Validity Scale (FBS), in detecting the over-reporting of PTSD symptoms in compensation seeking individuals during forensic evaluations (DeViva & Bloem, 2003; Franklin, Repasky, Thompson, Shelton, & Uddo, 2002; Frueh, Gold, & de Arellano, 1997; Greiffenstein, Baker, Axelrod, Peck, & Gervais, 2004; Rogers, Sewell, Martin, & Vitacco, 2003; Smith & Frueh, 1996).

The MMPI-2 Restructured Form (MMPI-2-RF; Ben-Porath & Tellegen, 2008) represents the latest version of the MMPI test and was developed in an effort to modernize the test by addressing the psychometric limitations of the MMPI and MMPI-2, as well as assessing personality and psychopathology in line with more contemporary theories (Ben-Porath & Tellegen, 2008). The focus of this paper will be the use of the MMPI-2-RF in the conceptualization of PTSD.

The MMPI-2-RF utilizes a hierarchical structure that incorporates both broadband and narrowly focused scales that measure personality and temperament, psychological symptoms, individual differences, and specific behavioral proclivities (Tellegen & Ben-Porath, 2008). The three Higher Order (H-O) scales represent broadband

measures of Emotional/Internalizing Dysfunction (EID), Thought Dysfunction (THD), and Behavioral/Externalizing Dysfunction (BXD), and provide clinicians with an overall evaluation of the general problem areas of the test-taker. These three broad dimensions of functioning represent consistent findings in larger meta-analytic and epidemiological investigations of psychopathology (e.g., Krueger & Markon, 2006). Moreover, they provide a dimensional representation of three commonly occurring code types in clinical settings (2-7/7-2; 4-9/9-4; 6-8/8-6; see Graham, 2006).

Following the H-O scales in the hierarchical structure of the MMPI-2-RF are the Restructured Clinical (RC) scales (Tellegen et al., 2003), which clarify the H-O scales with a more focused evaluation of symptom and trait patterns. In an effort to address the high inter-correlations between the Clinical Scales of the MMPI-2, which can complicate effective interpretation of multiple scale elevations, Tellegen et al. (2003) developed the RC scales prior to their inclusion in the MMPI-2-RF (Ben-Porath & Tellegen, 2008). These scales differ from the original Clinical Scales in that general emotional distress and dysfunction, which was identified as the main contributor to the high inter-correlations on the Clinical Scales on the MMPI-2 (Tellegen et al., 2003), is identified as a separate construct and found on the RC Demoralization scale (RCd). Demoralization is conceptually related to the general distress component found to be underlying many heterogeneous disorders (Brown, Chorpita, Barlow, 1998; Moses & Barlow, 2006). Indeed, measures of demoralization and negative affect have been found to correlate highly as indicators of psychological distress (Sellbom, Ben-Porath, & Bagby, 2008). Previous research has examined the use of the MMPI-2 RC scales in assessing constructs associated

with the symptom clusters of PTSD (Wolf et al., 2008). Specifically, Wolf and colleagues (2008) found that re-experiencing symptoms such as flashbacks were captured by Aberrant Experiences (RC8), emotional numbing was associated with Low Positive Emotions (RC2), and hyperarousal was related to Dysfunctional Negative Emotions (RC7). Additionally, somatic complaints and the overall emotional distress associated with the chronic and pervasive effects of the disorder were associated with Somatic Complaints (RC1) and Demoralization (RCd), respectively.

The most narrowly focused scales on the MMPI-2-RF are the Specific Problem (SP) scales, which are grouped in to indexes measuring somatic and cognitive complaints, internalizing and externalizing behaviors and cognitions, and interpersonal experiences. Several of the SP scales may also prove useful in case conceptualizations in assessing various symptoms of PTSD. Of the Interpersonal scales, the Family Problems scale (FML) should be considered, as the amount of post-trauma social support available to an individual can be a moderating factor on the development of pathology (Koenen, Stellman, Stellman, & Sommer, 2003; Ruscio, Ruscio, & Keane, 2002). Additionally, scores on Social Avoidance (SAV) and Disaffiliativeness (DSF) may provide insight into whether an individual is experiencing a feeling of detachment from others, consistent with the DSM-IV's description of avoidance symptoms (APA, 2000). Lastly, Substance Abuse (SUB) of the Externalizing Scales might capture alcohol and drug misuse, which has long been associated with a diagnosis of PTSD in various trauma populations (Bremner et al., 1996; Epstein et al., 1996). Please refer to Table 1 for a more information regarding MMPI-2-RF scales and PTSD symptoms.

Lastly, the MMPI-2 Personality Psychopathology Five (PSY-5) scales, developed by Harkness & McNulty

Table 1 *MMPI-2-RF and PTSD Symptoms*

MMPI-2-RF Scales				
PTSD Cluster/Symptom	H-O	RC	SP	PSY-5
<i>Re-Experiencing</i>	THD		AXY	PSYC-r
Recurrent distressing dreams of the event			AXY	
Intense psychological distress at exposure to cues reminiscent of the trauma			AXY	
Physiological reactivity at exposure to cues reminiscent of the trauma			AXY	
<i>Avoidance/Numbing</i>	EID	RC2, RC7		INTR-r
Avoidance of activities, places, or people associated with trauma			BRF	
Inability to recall an important aspect of the trauma			COG	
Diminished interest/participation in significant activities			SAV	
Feeling of detachment or estrangement from others			DSF	INTR-r
Restricted range of affect		RC2, RC7		INTR-r
Sense of a fore-shortened future		RC7	HLP	

Table 1 continued

<i>Hyperarousal</i>	BXD	RC7, RC9	AGGR-r, DISC-r
Sleep difficulty		MLS	
Irritability/anger		RC7 ANP,	DISC-r
Poor concentration		AGG	
Hyper-vigilance		COG	
Exaggerated startle response		RC6, RC9 RC1, RC9	AXY AXY

(1994) to measure dispositional abnormal personality characteristics, were revised for inclusion in the MMPI-2-RF. The PSY-5-r scales include Aggressiveness (AGGR-r), Psychoticism (PSYC-r), Disconstraint (DISC-r), Neuroticism/Negative Emotionality (NEGE-r), and Introversion/Low Positive Emotionality (INTR-r). Certain personality traits measured by the PSY-5 scales have been linked to PTSD symptom expression in recent research. Miller (2003) found that low levels of constraint and inhibition appear to act as moderating factors on the PTSD symptoms expressed in traumatized individuals. Additionally, Miller and colleagues (2004) utilized a cluster analysis of the PSY-5 scales in a sample of veterans and found three distinct clusters of PTSD symptom expression: a low-level pathology group, an externalizing group characterized by high scores on AGGR, NEGE, PSYC, and DISC, and an internalizing cluster characterized by high scores NEGE and high scores on INTR. This study replicated and extended findings from an earlier examination of posttraumatic response subtypes (Miller, Greif, & Smith, 2003), and similar findings have been found in samples of female

rape survivors (Miller & Resick, 2007) as well as workplace claimants (Sellbom & Bagby, 2009).

In addition to its various clinically substantive scale, the MMPI-2-RF contains a set of validity indicators that can be used to identify both non-content (e.g., random, acquiescent responding) and content-based response bias (e.g., over-reporting, under-reporting). The MMPI-2-RF over-reporting scales include revised versions of the Infrequency (F), Infrequency Psychopathology (Fp), and the Symptom Validity scale¹ (FBS). The Infrequent Somatic Responses (Fs; Wygant, Ben-Porath, & Arbisi, 2004) was added to the MMPI-2-RF and contains items infrequently endorsed in medical and chronic pain samples. The MMPI-2-RF over-reporting validity scales have been found to be effective markers of response bias in both civil/disability settings (Wygant, Ben-Porath, Arbisi, Berry, Freeman, & Heilbronner, 2009; Wygant, Anderson, Sellbom, Rapier, Allgeir, & Granacher, 2011) and criminal forensic settings (Sellbom, Toomey, Wygant, Kucharski, & Duncan, 2010). Additionally, the Response Bias Scale (RBS; Gervais, Ben-Porath, Wygant, & Green, 2007) was recently added to the MMPI-2-RF. This scale contains 28 items that were found to effectively discriminate between individuals who passed or failed cognitive symptom validity tests commonly used in forensic disability settings. The strength of these scales is their ability to detect various features of malingering, such as severe psychopathology (Fp-r) or overreported (Fp-r) or overreported neurocognitive impairment (RBS), and somatic exaggeration (Fs & FBS-r). As such, these validity scales lend themselves well to the assessment of disorders with hetero-

¹Previously labeled the Fake Bad Scale, this measure was re-named Symptom Validity to provide a more descriptive and less inferential label (Ben-Porath, Tellegen, & Graham, 2008).

geneous symptom constellations, such as PTSD, in both clinical settings and in forensic situations where the veracity of the symptom presentation must be examined.

Individual PTSD Case Conceptualizations Using the MMPI-2-RF

An MMPI-2-RF profile, completed as part of a disability evaluation with the third author for claims of psychological damage in the form of PTSD, suggest that the test has good clinical utility for case conceptualizations in individuals with the disorder.

The conceptualized case is based on a 43 year old divorced male who was involved in a vehicular collision that resulted in the death of the other driver. Immediately following the accident, the individual experienced a dissociative episode that lasted for approximately 36 hours. At the time of the assessment, which occurred 10 years following the initial trauma, the man reported experiencing occasional, brief dissociative episodes, typically triggered by odors he associated with the accident. His assessment resulted in a diagnosis of PTSD.

In terms of his MMPI-2-RF results, a review of his validity scales suggests that he was generally cooperative with the evaluation (see Figure 1). His non-content based validity scale elevation (i.e., VRIN-r, TRIN-r) were in the normal range and none of the over-reporting validity scales were indicative of symptom exaggeration. Although he perhaps attempted to portray himself in an overly virtuous light (L-r =76), he still endorsed problematic symptoms on a number of the substantive scales of the MMPI-2-RF. In the absence of any elevations on the substantive measures of the MMPI-2-RF, his score on L-r might reflect defensiveness; however, given that the remainder of his profile

shows marked elevations on scales reflecting the avoidance, re-experiencing, and hyperarousal symptom clusters of PTSD (APA, 2000), his score might actually reflect an orientation towards traditional values. Further supporting the veracity of his symptom report was his effort while testing and his non-elevated score on the Response Bias Scale. Indeed, his RBS t-score was 50 and he passed the three cognitive symptom validity tests administered during his evaluation, including 100% performance on the Word Memory Test and Medical Symptom Validity Test.




Figure 1. MMPI-2-RF Validity Scales. Excerpted from the MMPI-2-RF Manual for Administration, Scoring, and Interpretation by Yossef S. Ben-Porath and Auke Tellegen. Copyright © 2008 by the Regents of the University of Minnesota. Reproduced by permission of the University of Minnesota Press. All rights reserved. “Minnesota Multiphasic Personality Inventory-2-RF®” and “MMPI-2-RF®” are trademarks owned by the Regents of the University of Minnesota.

Legend: VRIN-r = Variable Response Inconsistency; TRIN-r = True Response Inconsistency; Fr= Infrequent Responses; Fp-r = Infrequent Psychopathology Responses; Fs = Infrequent Somatic Responses; FBS-r = Symptom Validity; L-r = Uncommon Virtues; K-r = Adjustment Validity.

This client presented scale elevations suggestive of both emotional and behavioral avoidance, which support a diagnosis of PTSD. His overall level of emotional dysfunction is suggested by his elevated score on the Emotional/Internalizing Dysfunction Higher-Order Scale (EID=68), which measures the dysfunctional affect, emotional numbing, and rumination.

Examination of the client's remaining profiles (Figures 2-4) provides more focused information about his symptom pattern, as well as his scores on the PSY-5-r scales (Figure 5), which suggests some potential underlying etiological factors to consider. His general sense of unhappiness, which may encompass the feelings of hopelessness that often occur as a result of PTSD, is exhibited by his elevation on Demoralization (RCd = 66). Hopelessness specifically appears to be a problem for this individual, as seen by his score on the Hopelessness/ Helplessness scale (HLP=69). Additionally, this individual had an elevated score on Negative Emotionality (NEGE-r=73) of the PSY-5-r scales. This PSY-5-r scale elevation is consistent with findings that identify negative emotionality as a consistent feature of both internalizing and externalizing subtypes of PTSD MMPI-2 profiles (Miller et al., 2004). Previous research has supported broad-band personality factors such as those measured by the PSY-5-r scales as providing etiological continuity between Axis I and Axis II disorders (Krueger, 2005), suggesting these personological variables play a key role in the development of pathology such as PTSD. In terms of behavioral avoidance (PTSD Criterion C), the most striking evidence for behavior change as a possible result of this individual's PTSD is found on the Behavior Restricting Fears scale (BRF=79). This score suggests a restriction of behavior and avoidance of normal activities as a result of intense fear or distress, which corre-

sponds with the DSM-IV symptom criteria for the disorder (APA, 2000). This score may indicate that the individual is determined to avoid recollections of the trauma by drastically decreasing the breadth of his activities, which allows




Figure 2. MMPI-2-RF Higher-Order (H-O) and Restructured Clinical (RC) Scales. Excerpted from the MMPI-2-RF Manual for Administration, Scoring, and Interpretation by Yossef S. Ben-Porath and Auke Tellegen. Copyright © 2008 by the Regents of the University of Minnesota. Reproduced by permission of the University of Minnesota Press. All rights reserved. “Minnesota Multiphasic Personality Inventory-2-RF®” and “MMPI-2-RF®” are trademarks owned by the Regents of the University of Minnesota.

Legend: EID = Emotional/Internalizing Dysfunction; THD = Thought Dysfunction; BXD = Behavioral/Externalizing Dysfunction; RCd = Demoralization; RC1 = Somatic Complaints; RC2 = Low Positive Emotions; RC3 = Cynicism; RC4 = Antisocial Behavior; RC6 = Ideas of Persecution; RC7 = Dysfunctional Negative Emotions; RC8 = Aberrant Experiences; RC9 = Hypomanic Activation.




Figure 3. MMPI-2-RF Somatic/Cognitive and Internalizing Scales. Excerpted from the MMPI-2-RF Manual for Administration, Scoring, and Interpretation by Yossef S. Ben-Porath and Auke Tellegen. Copyright © 2008 by the Regents of the University of Minnesota. Reproduced by permission of the University of Minnesota Press. All rights reserved. “Minnesota Multiphasic Personality Inventory-2-RF®” and “MMPI-2-RF®” are trademarks owned by the Regents of the University of Minnesota.

Legend: MLS = Malaise; GIC = Gastrointestinal Complaints; HPC = Head Pain Complaints; NUC = Neurological Complaints; COG = Cognitive Complaints; SUI = Suicidal/Death Ideation; HLP = Helplessness/Hopelessness; SFD = Self-Doubt; NFC = Inefficiency; STW = Stress/Worry; AXY = Anxiety; ANP = Anger Proneness; BRF = Behavior Restricting Fears; MSF = Multiple Specific Fears.

him to attenuate his trauma-induced fears in a negatively reinforcing pattern. Additionally, his score on the Social Avoidance Scale (SAV=70) suggests that he is avoiding people and social gatherings as well. This interpersonal avoidance is also characteristic of individuals experiencing




Figure 4. MMPI-2-RF Externalizing, Internalizing, and Interest Scales. Excerpted from the MMPI-2-RF Manual for Administration, Scoring, and Interpretation by Yossef S. Ben-Porath and Auke Tellegen. Copyright © 2008 by the Regents of the University of Minnesota. Reproduced by permission of the University of Minnesota Press. All rights reserved. “Minnesota Multiphasic Personality Inventory-2-RF®” and “MMPI-2-RF®” are trademarks owned by the Regents of the University of Minnesota.

Legend: JCP = Juvenile Conduct Problems; SUB = Substance Abuse; AGG = Aggression; ACT = Activation; FML = Family Problems; IPP = Interpersonal Passivity; SAV = Social Avoidance; SHY = Shyness; DSF = Disaffiliativeness; AES = Aesthetic-Literary Interests; MEC = Mechanical-Physical Interests.

PTSD (APA, 2000), and in this case may result from cynical beliefs stemming from the trauma that others are not to be trusted (RC3=70). Active interpersonal avoidance may also account for this individual’s score on the Shyness scale (SHY=75). While this individual may have originally had some shyness as a natural component of his




Figure 5. MMPI-2-RF PSY-5 Scales. Excerpted from the MMPI-2-RF Manual for Administration, Scoring, and Interpretation by Yossef S. Ben-Porath and Auke Tellegen. Copyright © 2008 by the Regents of the University of Minnesota. Reproduced by permission of the University of Minnesota Press. All rights reserved. “Minnesota Multiphasic Personality Inventory-2-RF®” and “MMPI-2-RF®” are trademarks owned by the Regents of the University of Minnesota.

Legend: AGGR-r = Aggressiveness-Revised; PSYC-r = Psychoticism-Revised; DISC-r = Disconstraint-Revised; NEGE-r = Negative Emotionality/Neuroticism-Revised; INTR-r = Introversion/Low Positive Emotionality-Revised

temperament, his elevation on SHY may reflect emotional disengagement from those around him since the accident.

Evidence of symptoms inherent to the re-experiencing symptom cluster (PTSD Criterion B) can also be discerned from the client’s MMPI-2-RF profile. His score on the Psychoticism scale of the PSY-5-r Scales is mildly elevated (PSYC-r=63). This score is suggestive of a

mild level of disconnect from reality. The intense psychological distress of reexperiencing a traumatic event in a vivid, recurrent manner may additionally elevate this scale, in addition to this individual’s particular style of dissociating in response to triggers associated with the original trauma. Moreover, Miller et al. (2004) suggested that Psychoticism might be measuring feelings of alienation and absorption, both of which are suggestive of PTSD.

Several scale elevations illustrate symptoms indicative of the Hyperarousal symptom cluster (PTSD Criterion D). In particular, the RC scale measuring Dysfunctional Negative Emotions is elevated on this profile (RC7=65). RC7 measures a wide range of dysfunctional negative emotions, such as maladaptive anxiety, anger, and irritability. Several SP scales provide further evidence of Criterion D symptoms. Indeed, his score on the Activation Scale (ACT=67) suggests a heightened excitation and energy level, which corresponds with the increased arousal that accompanies a diagnosis of PTSD. This maladaptive anxiety is also evidenced by elevations on Stress/Worry (STW=73) and Anxiety (AXY=91), as would be expected with an individual constantly reliving a traumatic event.

Discussion

PTSD is a complex disorder to assess in any setting, as a result of its heterogeneous symptom picture that is primarily discerned through self-report. The assessment of PTSD in a forensic setting further adds to the complexity of rendering a diagnosis, as the disorder is often a disability that is monetarily compensable. The MMPI-2-RF provides assistance in assessing empirically supported constructs associated with PTSD as a result of its hierar-

chical structure that differentiates internalizing and externalizing symptoms and its ability to isolate the impact of demoralization, or general emotional distress, which permeates many Axis I conditions. Additionally, the MMPI-2-RF contains validity scales that measure a variety of threats to protocol validity (e.g., exaggerated psychopathology, neurocognitive impairment, and physical health problems). These scales allow the clinician to form an opinion about the specific nature of potential symptom exaggeration, as PTSD can feigned in numerous ways (see Resnick et al., 2008 for a review).

The preceding case study and discussion should highlight the clinical utility of the MMPI-2-RF in interpreting individual presentations of PTSD. Further research, however, is needed to explore the association between the MMPI-2-RF and PTSD on a more empirical level.

References

- American Psychiatric Association. (1980). *Diagnostic and statistical manual of mental disorders* (3rd ed.). Washington, D.C.: American Psychiatric Association.
- American Psychiatric Association. (2000). *Diagnostic and statistical manual of mental disorders* (4th ed., Text revision). Washington, DC: American Psychiatric Association.
- Arbisi, P., Ben-Porath, Y., & McNulty, J. (2006). The ability of the MMPI-2 to detect feigned PTSD within the context of compensation seeking. *Psychological Services, 3*, 249-261.
- Archer, R., Buffington-Vollum, J., Stredny, R., & Handel, R. (2006). A survey of psychological test use patterns among forensic psychologists. *Journal of Personality Assessment, 87*, 84-94.
- Ben-Porath, Y.S., & Tellegen, A. (2008). *The Minnesota Multiphasic Personality Inventory -2 Restructured Form:*

- Manual for administration, scoring, and interpretation.* Minneapolis, MN: University of Minnesota Press.
- Ben-Porath, Y.S., Tellegen, A., & Graham, J.R. (2008). *The MMPI-2 Symptom Validity Scale.* Minneapolis, MN: University of Minnesota Press.
- Birmes, P., Hatton, L., Brunet, A., & Schmitt, L. (2003). Early historical literature for post-traumatic symptomatology. *Stress and Health 19*, 17-26.
- Blake, D. B., Weathers, F. W., Nagy, L. M., Kaloupek, D. G., Gusman, F. D., Charney, D. S., et al. (1995). The development of a Clinician-Administered PTSD Scale. *Journal of Traumatic Stress, 8*, 75-90.
- Borum, R. & Grisso, T. (1995). Psychological test use in criminal forensic evaluations. *Professional Psychology: Research and Practice, 26*, 465-473.
- Bremner, J.D., Southwick, S.M., Darnell, A., & Charney, D.S. (1996) Chronic PTSD in Vietnam combat veterans: Course of illness and substance abuse. *American Journal of Psychiatry, 153*, 369-375.
- Briere, J. (1995). *Trauma Symptom Inventory professional manual.* Odessa, FL: Psychological Assessment Resources.
- Brown, T., Chorpita, B., & Barlow, D. (1998). Structural relationships among dimensions of the DSM-IV anxiety and mood disorders and dimensions of negative affect, positive affect, and autonomic arousal. *Journal of Abnormal Psychology, 107*, 179-192.
- Brown, T. A., DiNardo, P. A., & Barlow, D. H. (1994). *Anxiety Disorders Interview Schedule for DSM-IV: Adult Version, Client Interview Schedule.* San Antonio, TX: The Psychological Corporation.
- Butcher, J.N., Graham, J.R., Ben-Porath, Y.S., Tellegen, A., Dahlstrom, W.G., & Kaemmer, B. (2001). *MMPI-2: Manual for the administration, scoring and interpretation.* Minneapolis: University of Minnesota Press.
- Calhoun, P., Earnst, K., Tucker, D., Kirby, A., & Beckham, J. (2000). Feigning combat-related posttraumatic stress disorder on the Personality Assessment Inventory. *Jour-*

- nal of Personality Assessment*, 75, 338-350.
- Camara, W., Nathan, J., & Puente, A. (2000). Psychological test usage: Implications in professional psychology. *Professional Psychology: Research and Practice*, 31, 141-154.
- Demakis, G., Gervais, R., & Rohling, M. (2008). The effect of failure on cognitive and psychological symptom validity tests in litigants with symptoms of post-traumatic stress disorder. *The Clinical Neuropsychologist*, 22, 879-895.
- DeViva, J., & Bloem, W. (2003). Symptom Exaggeration and Compensation Seeking Among Combat Veterans With Posttraumatic Stress Disorder. *Journal of Traumatic Stress*, 16, 503-507.
- Effendov, A., Sellbom, M., & Bagby, R. (2008). The utility and comparative incremental validity of the MMPI-2 and Trauma Symptom Inventory validity scales in the detection of feigned PTSD. *Psychological Assessment*, 20, 317-326.
- Elhai, J., Gray, M., Naifeh, J., Butcher, J., Davis, J., Falsetti, S., et al. (2005). Utility of the Trauma Symptom Inventory's Atypical Response Scale in Detecting Malingered Post-Traumatic Stress Disorder. *Assessment*, 12, 210-219.
- Elhai, J. D., Ruggiero, K. J., Frueh, B. C., Beckham, J. C., Gold, P. B., & Feldman, M. E. (2002). The Infrequency-Posttraumatic Stress Disorder Scale (Fptsd) for the MMPI-2: Development and initial validation with veterans presenting with combat-related PTSD. *Journal of Personality Assessment*, 79, 531-549.
- Epstein, J.N., Saunders, B.E., Kilpatrick, D.G., & Resnick, H.S. (1998). PTSD as a mediator between childhood rape and alcohol use in adult women. *Child Abuse and Neglect*, 22, 223-234.
- First, Michael B., Spitzer, Robert L, Gibbon Miriam, and Williams, Janet B.W.: Structured Clinical Interview for DSM-IV-TR Axis I Disorders, Research Version,

- Patient Edition (SCID-I/P). New York: Biometrics Research, New York State Psychiatric Institute, November 2002.
- Foa, E., Cashman, L., Jaycox, L., & Perry, K. (1997). The validation of a self-report measure of PTSD: The Posttraumatic Diagnostic Scale. *Psychological Assessment*, 9, 445-451.
- Foa, E., Hembree, E., & Rothbaum, B. (2007). *Prolonged exposure therapy for PTSD: Emotional processing of traumatic experiences: Therapist guide*. New York, NY US: Oxford University Press.
- Franklin, C., Repasky, S., Thompson, K., Shelton, S., & Uddo, M. (2003). Assessment of response style in combat veterans seeking compensation for posttraumatic stress disorder. *Journal of Traumatic Stress*, 16, 251-255.
- Franklin, C., Repasky, S., Thompson, K., Shelton, S., & Uddo, M. (2002). Differentiating over-reporting and extreme distress: MMPI-2 use with compensation-seeking veterans with PTSD. *Journal of Personality Assessment*, 79, 274-285.
- Franklin, C.L. & Thompson, K.E. (2005). Response style and Posttraumatic Stress Disorder (PTSD): A review. *Journal of Trauma and Dissociation*, 6, 105-123.
- Frueh, B., Elhai, J., Gold, P., Monnier, J., Magruder, K., Keane, T., et al. (2003). Disability compensation seeking among veterans evaluated for posttraumatic stress disorder. *Psychiatric Services*, 54, 84-91.
- Frueh, B., Gold, P., & de Arellano, M. (1997). Symptom over-reporting in combat veterans evaluated for PTSD: Differentiation on the basis of compensation seeking status. *Journal of Personality Assessment*, 68, 369-384.
- Frueh, B., Smith, D., & Barker, S. (1996). Compensation seeking status and psychometric assessment of combat veterans seeking treatment for PTSD. *Journal of Traumatic Stress*, 9, 427-440.
- Gervais, R.O., Ben-Porath, Y.S., Wygant, D.B., Green, P.

- (2007). Development and validation of a Response Bias Scale (RBS) for the MMPI-2. *Assessment, 14*, 196-208.
- Graham, J.R. (2006). *MMPI-2: Assessing personality and psychopathology*. New York: Oxford University Press.
- Greenberg, S., Otto, R., & Long, A. (2003). The utility of psychological testing in assessing emotional damages in personal injury litigation. *Assessment, 10*, 411-419.
- Greiffenstein, M.F., Baker, W.J., Axelrod, B., Peck, E.A., & Gervais, R. (2004). The Fake Bad Scale and MMPI-2 F-family in detection of implausible psychological trauma claims. *The Clinical Neuropsychologist, 18*, 573-590.
- Guriel-Tennant, J., & Fremouw, W. (2006). Impact of trauma history and coaching on malingering of posttraumatic stress disorder using the PAI, TSI, and M-FAST. *Journal of Forensic Psychiatry & Psychology, 17*, 577-592.
- Haisch, D.C. & Meyers, L.S. (2004). MMPI-2 assessed post-traumatic stress disorder related to job stress, coping, and personality in police agencies. *Stress and Health, 20*, 223-229.
- Harkness, A., & McNulty, J. (1994). The Personality Psychopathology Five (PSY-5): Issues from the pages of a diagnostic manual instead of a dictionary. *Differentiating normal and abnormal personality* (pp. 291-315). New York, NY US: Springer Publishing Co.
- Harkness, A., McNulty, J., & Ben-Porath, Y. (1995). The Personality Psychopathology Five (PSY-5): Constructs and MMPI-2 scales. *Psychological Assessment, 7*, 104-114.
- Hathaway, S.R. & McKinley, J.C. (1943). *The Minnesota Multiphasic Personality Inventory*. Minneapolis, MN: University of Minnesota Press.
- Hawes, S., & Boccaccini, M. (2009). Detection of over-reporting of psychopathology on the Personality Assessment Inventory: A meta-analytic review. *Psychological Assessment, 21*, 112-124.

- Helmes, E., & Reddon, J. (1993). A perspective on developments in assessing psychopathology: A critical review of the MMPI and MMPI-2. *Psychological Bulletin, 113*, 453-471.
- Keane, T. M., Caddell, J. M., & Taylor, K. L. (1988). Mississippi Scale for Combat Related Posttraumatic Stress Disorder: Three studies in reliability and validity. *Journal of Consulting and Clinical Psychology, 56*, 85-90
- Keane, T. M., Malloy, P. F., & Fairbank, J. A. (1984). Empirical development of an MMPI subscale for the assessment of combat-related posttraumatic stress disorder. *Journal of Consulting and Clinical Psychology, 52*, 888-891.
- Koenen, K.C., Stellman, J.M., Stellman, S.D., Sommer, J.F. (2003). Risk factors for course of posttraumatic stress disorder among Vietnam veterans: A 14-year follow-up of American Legionnaires. *Journal of Consulting and Clinical Psychology, 71*, 980-986.
- Krueger, R.F. (2005). Continuity of Axes I and II: Toward a unified model of personality, personality disorders, and clinical disorders. *Journal of Personality Disorders, 19*, 233-261.
- Krueger, R. F., & Markon, E. (2006). Reinterpreting comorbidity: A model-based approach to understanding and classifying psychopathology. *Annual Review of Clinical Psychology, 2*, 111-133.
- Lasiuk, G., & Hegadoren, K. (2006). Posttraumatic Stress Disorder Part I: Historical Development of the Concept. *Perspectives in Psychiatric Care, 42*(1), 13-20.
- Lees-Haley, P.R. (1992). Psychodiagnostic test usage by forensic psychologists. *American Journal of Forensic Psychology, 10*, 25-30.
- McKeever, V., & Huff, M. (2003). A diathesis-stress model of posttraumatic stress disorder: Ecological, biological, and residual stress pathways. *Review of General Psychology, 7*, 237-250.
- Miller, H., Goldberg, J., & Streiner, D. (1995). What's in a

- name? The MMPI-2 PTSD scales. *Journal of Clinical Psychology*, 51, 626-631.
- Miller, M. W. (2003). Personality and the etiology and expression of PTSD: A three-factor model perspective. *Clinical Psychology: Science and Practice*, 10, 373-393.
- Miller, M., Greif, J., & Smith, A. (2003). Multidimensional Personality Questionnaire profiles of veterans with traumatic combat exposure: Externalizing and internalizing subtypes. *Psychological Assessment*, 15, 205-215.
- Miller, M.W., Kaloupek, D.G., Dillon, A.L., & Keane, T.M. (2004). Externalizing and internalizing subtypes of combat-related PTSD: A replication and extension using the PSY-5 scales. *Journal of Abnormal Psychology*, 113, 636-645.
- Miller, M., & Resick, P. (2007). Internalizing and externalizing subtypes in female sexual assault survivors: Implications for the understanding of complex PTSD. *Behavior Therapy*, 38, 58-71.
- Minneka, S., Watson, D., & Clark, L.A. (1998). Comorbidity of anxiety and unipolar mood disorders. *Annual Review of Psychopathology*, 49, 377-412.
- Mollica, R. F., Caspi-Yavin, Y., Bollini, P., Truong, T., Tor, S., & Lavelle, J. (1992). The Harvard Trauma Questionnaire: Validating a cross-cultural instrument for measuring torture, trauma, and posttraumatic stress disorder in Indochinese refugees. *The Journal of Nervous and Mental Disease*, 180, 111-116.
- Morey, L. C. (1991). *Personality Assessment Inventory: Professional manual*. Tampa, FL: Psychological Assessment Resources.
- National Center for Posttraumatic Stress Disorder (2007). *Fact sheet: Epidemiological facts about PTSD; What causes Posttraumatic Stress Disorder? How common is it? Who gets it?* http://ncptsd.va.gov/ncmain/ncdocs/fact_shts/fs_epidemiological.html.

- Ozer, E., & Weiss, D. (2004). Who develops posttraumatic stress disorder?. *Current Directions in Psychological Science*, 13, 169-172.
- Resnick, P.J., West, S., & Payne, J.W. (2008). Malingering of posttraumatic stress disorder. In R. Rogers (ed.) *Clinical Assessment of Malingering and Deception* (3rd ed.). New York: The Guilford Press.
- Rogers, R., Sewell, K. W., Martin, M. A., & Vitacco, M. J. (2003). Detection of feigned mental disorders: A meta-analysis of the MMPI-2 and malingering. *Assessment*, 10, 160-177.
- Rubenzler, S. (2009). Posttraumatic stress disorder: Assessing response style and malingering. *Psychological Injury and Law*, 2, 114-142.
- Ruscio, A., Ruscio, J., & Keane, T. (2002). The latent structure of posttraumatic stress disorder: A taxometric investigation of reactions to extreme stress. *Journal of Abnormal Psychology*, 111, 290-301.
- Sellbom, M. & Bagby, R.M. (2009). Identifying PTSD personality subtypes in a workplace trauma sample. *Journal of Traumatic Stress*, 22, 471-475.
- Sellbom, M., Toomey, J. A., Wygant, D. B., Kucharski, L. T., & Duncan, S. (2010). *Psychological Assessment*, 22, 22-31.
- Schnurr, P., Friedman, M. J., & Bernardy, N. C. (2002). Research on posttraumatic stress disorder: Epidemiology, pathophysiology, and assessment. *Journal of Clinical Psychology*, 58, 877-889.
- Schnurr, P., Lunney, C., & Sengupta, A. (2004). Risk factors for the development versus maintenance of posttraumatic stress disorder. *Journal of Traumatic Stress*, 17, 85-95.
- Slick, D. J., Sherman, E. M., & Iverson, G. L. (1999). Diagnostic criteria for malingered neurocognitive dysfunction: Proposed standards for clinical practice and research. *The Clinical Neuropsychologist*, 13, 545-561.
- Smith, D., & Frueh, B. (1996). Compensation seeking, comor-

- bidity, and apparent exaggeration of PTSD symptoms among Vietnam combat veterans. *Psychological Assessment*, 8, 3-6.
- Souza, W. F., Figueira, I., Mendlowicz, M. V., Volchan, E., Mendonça-de-Souza, A. C., Duarte, A. F. A., Monteiro da Silva, Ângela M., Marques-Portella, C., Mari, J. d. J., & Coutinho, E. S. F. (2008). Negative affect predicts posttraumatic stress symptoms in Brazilian volunteer United Nations Peacekeepers in Haiti. *Journal of Nervous and Mental Disease*, 196, 852-855.
- Tellegen, A., & Ben-Porath, Y.S. (2008). *The Minnesota Multiphasic Personality Inventory -2 Restructured Form: Technical manual*. Minneapolis, MN: University of Minnesota Press.
- Watson, D. (2005). Rethinking the mood and anxiety disorders: A quantitative hierarchical model for DSM-V. *Journal of Abnormal Psychology*, 114, 522-536.
- Wolf, E.J., Miller, M.W., Orazem, R.J., Weierich, M.R., Castillo, D.T., Millford, J., Kaloupek, D.G., & Keane, T.M. (2008). The MMPI-2 Restructured Clinical scales in the assessment of posttraumatic stress disorder and comorbid disorders. *Psychological Assessment*, 20, 327-340.
- Wygant, D.B., Anderson, J.L., Sellbom, M., Rapier, J.L., Allgeier, L.M., & Granacher, R.P. (2011). Association of the MMPI-2 Restructured Form (MMPI-2-RF) Validity Scales with Structured Malingering Criteria. *Psychological Injury and Law*, 4, 13-23.
- Wygant, D.B., Ben-Porath, Y.S., & Arbisi, P.A. (May, 2004). *Development and initial validation of a scale to detect somatic over-reporting*. Paper presented at the 39th Annual Symposium of Recent Developments in the Use of the MMPI-2 and the MMPI-A, Minneapolis, MN.
- Wygant, D.B., Ben-Porath, Y.S., Arbisi, P.A., Berry, D.T.R., Freeman, D.B., & Heilbronner, R.L. (2009). Examination of the MMPI-2 Restructured Form (MMPI-2-RF) Validity Scales in civil forensic settings: Findings from

simulation and known group samples. *Archives of Clinical Neuropsychology*, 24, 671-680.