

Eastern Kentucky University

Encompass

EKU Faculty and Staff Scholarship

Faculty and Staff Scholarship Collection

3-2008

Using E. Coli and Bacteroides Distribution and Abundance in a Eutrophic Lake as a Tracer for Nutrient Inputs, Wilgreen Lake, Madison County, Kentucky

Theresa A. Aguiar

Walter S. Borowski

Eastern Kentucky University, w.borowski@eku.edu

Alice C. Layton

University of Tennessee, Knoxville

Larry McKay

University of Tennessee, Knoxville

Follow this and additional works at: https://encompass.eku.edu/fs_research



Part of the [Biogeochemistry Commons](#), [Environmental Health and Protection Commons](#), [Environmental Indicators and Impact Assessment Commons](#), and the [Environmental Monitoring Commons](#)

Recommended Citation

Aguiar, T.A., W.S. Borowski, A.C. Layton, L. McKay, 2008. Using E. coli and Bacteroides distribution and abundance in a eutrophic lake as a tracer for nutrient inputs, Wilgreen Lake, Madison County, Kentucky. GSA Abstracts with Programs, Vol. 40, No. 5.

This Conference Presentation is brought to you for free and open access by the Faculty and Staff Scholarship Collection at Encompass. It has been accepted for inclusion in EKU Faculty and Staff Scholarship by an authorized administrator of Encompass. For more information, please contact Linda.Sizemore@eku.edu.

Using *E. coli* and *Bacteroides* distribution and abundance in a eutrophic lake as a tracer for nutrient inputs, Wilgreen Lake, Madison County, Kentucky

Theresa A. Aguiar and Walter S. Borowski

Department of Geography and Geology
Eastern Kentucky University
Richmond, KY 40475
(theresa_aguiar@eku.edu; w.borowski@eku.edu)

Alice C. Layton and Larry McKay

University of Tennessee
Knoxville, TN 37996

Wilgreen Lake is a eutrophic lake that has been listed on the EPA's 303d list as nutrient impaired. Potential sources of this impairment are likely from humans, cattle manure, and fertilizers. We suspect that the majority of nutrients originate from human sources, namely from septic tank effluent emanating from key housing developments ringing the lakeshore. We test our hypothesis with conventional microbial assays (*Escherichia coli*) and RT-PCR techniques (*Bacteroides*).

We took water samples at 19 sampling locations on 4 occasions, and measured the abundance of *Escherichia coli* using IDEXX methods. Corresponding sub-samples slated for potential PCR analysis were stored at -40°C. We chose PCR assay candidates on the basis of elevated *E. coli* levels, and the probability of differing source contributions.

There is a systematic decline in *E. coli* microbial abundance distal to developments with closely-spaced septic units. This suggests that the principal source of microbial input is from septic systems; however, we cannot eliminate the possibility that fecal microbes are introduced into the lake via inflows.

We used quantitative PCR analysis to measure *Bacteroides* abundance, and to distinguish between human and cattle sources. We measured 14 samples and found total fecal microbe concentrations in all samples targeting all *Bacteroides* species ranged from 45 mg/L to 142 mg/L. Unlike other studies, there was no apparent relationship between the concentration of all *Bacteroides* species and that of *E. coli*.

We also attempted to quantitatively determine the proportion of *Bacteroides* contributions from specific sources, namely human and bovine fecal matter. Although fecal contamination was measured in all 14 samples, only 1 sample had significant amounts of human fecal contamination (21%) as measured by the human-associated *Bacteroides* assay. None of the samples had significant amounts bovine fecal concentration as measured by the bovine-associated *Bacteroides* assay. These inconclusive results suggest that either there are other unidentified sources of fecal contamination by *Bacteroides* and/or *E. coli*, or that the prevailing drought conditions skewed our results by not capturing fecal transport effects due to lack of surface and/or groundwater flow.