


4-2007

Dynamics of a eutrophic lake (Wilgreen Lake, Madison County, Kentucky): A first step in cleansing a lake system impaired by nutrient loading

Erin C. Jolly
Eastern Kentucky University

Walter S. Borowski
Eastern Kentucky University

Follow this and additional works at: https://encompass.eku.edu/fs_research

 Part of the [Environmental Indicators and Impact Assessment Commons](#), [Environmental Monitoring Commons](#), and the [Fresh Water Studies Commons](#)

Recommended Citation

Jolly, E.C., W.S. Borowski, 2007. Dynamics of a eutrophic lake (Wilgreen Lake, Madison County, Kentucky): A first step in cleansing a lake system impaired by nutrient loading. EKU Undergraduate Presentation Showcase, 13 April 2007.

This Poster is brought to you for free and open access by Encompass. It has been accepted for inclusion in EKU Faculty and Staff Scholarship by an authorized administrator of Encompass. For more information, please contact Linda.Sizemore@eku.edu.

**Dynamics of a eutrophic lake (Wilgreen Lake, Madison County, Kentucky):
A first step in cleansing a lake system impaired by nutrient loading**

Erin C. Jolly and Walter S. Borowski

Department of Earth Sciences and ECU Environmental Research Institute
Eastern Kentucky University
Richmond KY 40475-3102

Wilgreen Lake (Madison County, Kentucky) is a eutrophic lake formed by damming Taylor Fork, part of the Silver Creek watershed. The lake is listed “nutrient impaired” by the Commonwealth and the EPA, and it is likely that nutrient input from human activities is affecting water quality. Our study aims first to characterize the physical characteristics and water quality of the lake (2006), and then to determine the specific proportion of nutrient inputs (2007) to the lake with the ultimate aim of improving its water quality.

Research started in May 2006 with work occurring throughout the 2006 field season with the intent of establishing a baseline for key lake parameters. We used a YSI probe to measure temperature, conductivity, oxygen concentration, and pH, and assayed for total ammonia nitrogen using the sodium hypochlorite, colorimetric method. The lake was already strongly stratified in May with disoxic and anoxic water below about 4 meters. Stratification strengthened in the summer with the disoxic-oxic boundary moving upward to about 3 meters, showing a sharper gradient oxygen gradient. Ammonium concentrations are typically zero in the oxic zone, and increase in concentration with increasing water depth in anoxic waters to about 5 ppm.

EKU Undergraduate Presentation Showcase, 13 April 2007.