

January 2018

Positive Symmetric Solutions Of A Boundary Value Problem With Dirichlet Boundary Conditions

Tek Nath Dhakal
Eastern Kentucky University

Follow this and additional works at: <https://encompass.eku.edu/etd>



Part of the [Analysis Commons](#)

Recommended Citation

Dhakal, Tek Nath, "Positive Symmetric Solutions Of A Boundary Value Problem With Dirichlet Boundary Conditions" (2018). *Online Theses and Dissertations*. 563.
<https://encompass.eku.edu/etd/563>

This Open Access Thesis is brought to you for free and open access by the Student Scholarship at Encompass. It has been accepted for inclusion in Online Theses and Dissertations by an authorized administrator of Encompass. For more information, please contact Linda.Sizemore@eku.edu.

POSITIVE SYMMETRIC SOLUTIONS OF A BOUNDARY VALUE
PROBLEM WITH DIRICHLET BOUNDARY CONDITIONS


BY

TEK NATH DHAKAL

THESIS APPROVED:




Chair, Advisory Committee



Member, Advisory Committee



Member, Advisory Committee



Dean, Graduate School

STATEMENT OF PERMISSION TO USE

In presenting this thesis in partial fulfillment of the requirements for a M.S. degree at Eastern Kentucky University, I agree that the Library shall make it available to borrowers under rules of the Library. Brief quotations from this thesis are allowable without special permission, provided that accurate acknowledgment of the source is made. Permission for extensive quotation from or reproduction of this thesis may be granted by my major professor, or, in his absence, by the Head of Interlibrary Services when, in the opinion of either, the proposed use of the material is for scholarly purposes. Any copying or use of the material in this thesis for financial gain shall not be allowed without my written permission.

Signature 

Date 11/06/2018

POSITIVE SYMMETRIC SOLUTIONS OF A BOUNDARY VALUE PROBLEM
WITH DIRICHLET BOUNDARY CONDITIONS

BY

TEK N. DHAKAL

Submitted to the Faculty of the Graduate School of
Eastern Kentucky University
in partial fulfillment of the requirements for the degree of

MASTER OF SCIENCE

2018

© Copyright by TEK N. DHAKAL, 2018
All Rights Reserved.

DEDICATION

This thesis is dedicated to my parents.

ACKNOWLEDGEMENTS

I would like to take this opportunity to thank my thesis advisor, Dr. Jeffrey Neugebauer, for his honest advice and availability. Without his support and encouragement, this thesis would not be possible. I would also like to thank my thesis committee members, Dr. Steve Szabo and Dr. Atilla Sit, for their guidance.

ABSTRACT

We apply a recent extension of a compression-expansion fixed point theorem of function type to a second order boundary value problem with Dirichlet boundary conditions. We show the existence of positive symmetric solutions of this boundary value problem.

TABLE OF CONTENTS

CHAPTER	PAGE
1 INTRODUCTION	1
2 THE FIXED POINT THEOREM	5
2.1 Definitions	5
2.2 The Fixed Point Theorem	5
3 POSITIVE SYMMETRIC SOLUTIONS	7
3.1 Preliminaries	7
3.2 Positive Symmetric Solutions	8
3.3 Example	16
BIBLIOGRAPHY	20

1 INTRODUCTION

There have been many studies in the past on finding positive solutions of boundary value problems. In 1979, Leggett and Williams [14] came up with fixed point theorems which guaranteed the existence of solutions of the boundary value problems. The first theorem helps find a fixed point, the second one guarantees the existence of at least two fixed points, and the other two theorems guarantee the existence of at least three fixed points. The following theorem is the original Leggett-Williams theorem which guarantees the existence of a fixed point.

Theorem 1.1 (Leggett-Williams [14]). *Consider a Banach space B , the cone K , and the operator T . For α a positive concave functional, we define the following subsets of K :*

$$K_c = \{x \in K : \|x\| < c\},$$

$$K(\alpha, b, d) = \{x \in K : b \leq \alpha(x), \|x\| \leq d\}, \text{ and}$$

Suppose that $A : K_c \rightarrow K$ is completely continuous and suppose there exist a concave positive functional α with $\alpha(x) \leq \|x\|$ ($x \in K$) and numbers $b > a > 0$ ($b \leq c$) satisfying the following conditions:

(1) $\{x \in S(\alpha, a, b) : \alpha(x) > a\} \neq \emptyset$, and $\alpha(Ax) > a$ if $x \in S(\alpha, a, b)$;

(2) $Ax \in K_c$ if $x \in S(\alpha, a, c)$; and

(3) $\alpha(Ax) > a$ for all $x \in S(\alpha, a, c)$ with $\|Ax\| > b$.

Then A has a fixed point x in $S(\alpha, a, c)$.

Avery and other co-authors have generalized the Leggett-Williams theorem over the years, modifying the conditions, and have proven their theorem guarantees the existence of a positive solution. Some recent works of Avery, Anderson, and Henderson [10, 13] have also proven that there exists at least one positive solution for a right focal

boundary value problem. The compression-expansion fixed point theorem [10] helps find the fixed point when there is nonnegative continuous concave functionals and nonnegative continuous convex functions on \mathcal{P} , and when operator T is completely continuous. Some of the applications of the extensions of the Leggett-Williams fixed point theorem and the compression-expansion fixed point theorem are stated below.

In 2012, Altwaty and Eloë [2] have also used an extension of the original Leggett-Williams fixed point theorem to show the existence of at least one positive solution of a two point boundary value problem of the k th order differential equation.

In 2012, Avery, Eloë, and Henderson [8], used an extension of Leggett-Williams type fixed point theorem to study a two point boundary value problem for an ordinary differential equation that has a fourth order

$$x^{(iv)}(t) = f(x(t)), \quad t \in [0, 1],$$

$$x^{(i)}(0) = 0, \quad x^{(i)}(1) = 0, \quad i = 0, 1.$$

Similarly, in 2013, Altwaty and Eloë [1] used an extension of a Leggett-Williams in order to show that operator has at least one positive solution. They worked on two point conjugate boundary value problem

$$(-1)^n x^{(2n)}(t) = f(x(t)), \quad t \in [0, 1],$$

$$x^{(i)}(0) = 0, \quad x^{(i)}(1) = 0, \quad i = 0, 1, \dots, n$$

where n is a positive integer.

In 2016, Avery, Anderson, and Henderson [10] provided a modified version of the fixed point theorem which is known as the extension of the compression-expansion fixed point theorem. They used a compression-expansion fixed point theorem to show

the existence of a positive solution for a right focal boundary value problem

$$x''(t) + f(x(t)) = 0, \quad t \in (0, 1),$$

$$x(0) = 0 = x'(1),$$

where $f : \mathbb{R} \rightarrow [0, \infty)$ is continuous. If x is a fixed point of the operator T defined by

$$Tx(t) := \int_0^1 G(t, s)f(x(s))ds,$$

where $G(t, s) = \min\{t, s\}$, $(t, s) \in [0, 1] \times [0, 1]$, then x is a solution of the boundary value problem mentioned above.

In this paper, we will show the existence of a positive symmetric solution of the boundary value problem by using a compression-expansion fixed point theorem of function type [10]. Here, we have a second order boundary value problem with Dirichlet boundary value conditions. We will apply the fixed point theorem in order to show the existence of a positive symmetric solution to the differential equation

$$x''(t) + f(x(t)) = 0, \quad t \in (0, 1), \tag{1.1}$$

when it has the Dirichlet boundary conditions

$$x(0) = 0 = x(1), \tag{1.2}$$

where $f : \mathbb{R} \rightarrow [0, \infty)$ is continuous.

In Chapter 2, we have definitions and fixed point theorem. Under the fixed point theorem, we have extension of compression-expansion fixed point theorem. In Chapter 3, we have application of the extension of compression-expansion fixed point theorem and preliminaries. In addition to that, we have the properties of the Green's function

that we use throughout the paper, and the theorem we came up with. There are different conditions presented in the theorem, and if all these conditions are satisfied then we will show that our operator T has a fixed point $x^* \in A(\beta, b, \alpha, a)$. In other words, we will show the existence of a positive symmetric solution of (1.1), (1.2). We are motivated to work on this problem because of some of the recent works of Avery, Anderson, and Henderson [10, 13]. For more examples of Avery type fixed point theorems, see [3, 4, 5, 6, 7, 9, 11, 12].

2 THE FIXED POINT THEOREM

2.1 Definitions

Let us begin with some basic definitions.

Definition 1. A complete normed vector space is called a *Banach space*.

Definition 2. Let E be a real Banach space. A nonempty closed convex set $\mathcal{P} \subset E$ is called a *cone* if for all $x \in \mathcal{P}$ and $\lambda \geq 0$, $\lambda x \in \mathcal{P}$, and if $x, -x \in \mathcal{P}$ then $x = 0$.

Definition 3. A map α is said to be a *nonnegative continuous concave functional* on a cone \mathcal{P} of a real Banach space E if $\alpha : \mathcal{P} \rightarrow [0, \infty)$ is continuous and if

$$\alpha(tx + (1 - t)y) \geq t\alpha(x) + (1 - t)\alpha(y)$$

for all $x, y \in \mathcal{P}$ and $t \in [0, 1]$. Similarly, we say the map β is a *nonnegative continuous convex functional* on a cone \mathcal{P} of a real Banach space E if $\beta : \mathcal{P} \rightarrow [0, \infty)$ is continuous and if

$$\beta(tx + (1 - t)y) \leq t\beta(x) + (1 - t)\beta(y)$$

for all $x, y \in \mathcal{P}$ and $t \in [0, 1]$.

Definition 4. An operator is called *completely continuous* if it is continuous and maps bounded sets into sets whose closures are compact.

2.2 The Fixed Point Theorem

Definition 5. Let A be a relatively open subset of a cone \mathcal{P} , b and c positive real numbers, α a concave functional on \mathcal{P} , and β a convex functional on \mathcal{P} . We say that the open set

$$\mathcal{P}(\beta, b) = \{x \in \mathcal{P} : \beta(x) < b\}$$

is a *functional wedge* on a cone, that a set of the form

$$\mathcal{P}(\beta, b, \alpha, c) = \{x \in \mathcal{P} : c < \alpha(x) \text{ and } \beta(x) < b\}$$

is a *functional frustum* of a cone, and if A is an open subset of the cone \mathcal{P} , we say that

$$A(\beta, b, \alpha, c) = \{x \in A : c < \alpha(x) \text{ and } \beta(x) < b\}$$

is an interval of functional type.

Theorem 2.1 (Extension of Compression-Expansion Fixed Point Theorem [10]).

Suppose \mathcal{P} is a cone in a real Banach space E , A is a relatively open subset of \mathcal{P} , α and ψ are nonnegative continuous concave functionals on \mathcal{P} , β and θ are nonnegative continuous convex functionals on \mathcal{P} , and $T : \mathcal{P} \rightarrow \mathcal{P}$ is a completely continuous operator. If there exist nonnegative numbers a, b, c , and d such that

(A1) $A(\beta, b, \alpha, a)$ is bounded, $A(\beta, b, \alpha, a) \cap A(\theta, c, \psi, d) \neq \emptyset$, and if $x \in \partial A \cap \overline{\mathcal{P}(\beta, b, \alpha, a)} = \emptyset$ then $Tx \neq x$;

(A2) if $x \in \partial A(\beta, b, \alpha, a)$ with $\alpha(x) = a$ and either $\theta(x) \leq c$ or $\theta(Tx) > c$, then $\alpha(Tx) > a$;

(A3) if $x \in \partial A(\beta, b, \alpha, a)$ with $\beta(x) = b$ and either $\psi(Tx) < d$ or $\psi(x) \geq d$, then $\beta(Tx) < b$; then T has a fixed point $x^* \in A(\beta, b, \alpha, a)$.

3 POSITIVE SYMMETRIC SOLUTIONS

3.1 Preliminaries

Define the Banach space

$$E = C^{(1)}[0, 1] = \{x : [0, 1] \rightarrow \mathbb{R} : x' \text{ is continuous}\}$$

with norm

$$\|x\| = \max_{t \in [0, 1]} |x(t)| + \max_{t \in [0, 1]} |x'(t)|.$$

Define the cone $\mathcal{P} \subseteq E$ by

$$\mathcal{P} = \{x \in E : x(1-t) = x(t), x \text{ is nondecreasing on } [0, 1/2], \\ \text{nonnegative on } [0, 1], \text{ and concave on } [0, 1]\}.$$

The Green's function for $-x'' = 0$, (1.2), is given by

$$G(t, s) = \begin{cases} t(1-s), & 0 < t \leq s \leq 1, \\ s(1-t), & 0 \leq s \leq t \leq 1. \end{cases}$$

Therefore, x is a solution of (1.1), (1.2) if and only if x solves the integral equation

$$x(t) = \int_0^1 G(t, s) f(x(s)) ds, \quad t \in [0, 1].$$

Notice the derivative of the Green's function with respect to t is given by

$$\frac{\partial}{\partial t} G(t, s) = \begin{cases} (1-s), & 0 < t \leq s \leq 1, \\ -s, & 0 \leq s \leq t \leq 1. \end{cases}$$

Lemma 3.1. *The function G satisfies the following properties.*

1. $G(t, s) \in C([0, 1] \times [0, 1])$ with $G(t, s) \geq 0$ for all $(t, s) \in [0, 1] \times [0, 1]$.
2. $G(1 - t, 1 - s) = G(t, s)$ for all $(t, s) \in [0, 1] \times [0, 1]$.
3. For any $y, w \in [0, 1/2]$ with $y \leq w$,

$$yG(w, s) \leq wG(y, s).$$

Define the operator $T : E \rightarrow E$ by

$$Tx(t) = \int_0^1 G(t, s)f(x(s))ds, \quad t \in [0, 1].$$

If x is a fixed point of T , i.e. if $Tx = x$, then x solves (1.1), (1.2). In [2], it is shown that $T : \mathcal{P} \rightarrow \mathcal{P}$ and that T is completely continuous.

Define the nonnegative continuous functionals by

$$\beta(x) = \max_{t \in [0, 1/2]} x(t) = x(1/2),$$

$$\psi(x) = x'(1/8),$$

and

$$\alpha(x) = \min_{t \in [1/8, 1/2]} x(t) = x(1/8).$$

3.2 Positive Symmetric Solutions

Theorem 3.2. *Choose k_1, k_2, k_3, k_4 , and k_5 positive so that*

$$(a) \quad \frac{128}{7}k_4 < 8k_3,$$

$$(b) \quad k_1 < \left(\frac{83}{40}\right) \left(k_3 - \frac{9}{256}k_2\right),$$

$$(c) \quad 16 \left(k_4 - \left(\frac{192}{913}\right)k_1\right) < k_3 - k_5,$$

$$(d) \frac{4}{55} \left(k_3 - \left(\frac{9}{256} \right) k_2 \right) > k_4 - \left(\frac{192}{913} \right) k_1,$$

$$(e) k_2 > 64 \left(k_4 - \left(\frac{192}{913} \right) k_1 \right),$$

$$(f) k_2 > \left(\frac{2048}{913} \right) k_1,$$

$$(g) k_3 - k_5 > \left(\frac{512}{913} \right) k_1,$$

$$(h) k_3 > \frac{1}{8} (k_5 + 8k_1), \text{ and}$$

$$(i) k_4 < \left(\frac{1}{8} \right) (k_5 + 8k_1).$$

Let $f : [0, \infty) \rightarrow [0, \infty)$ be a continuous function and b be a positive real number such that

$$(H1) f(w) > \frac{4,096}{913} k_1 b \text{ for } w \in [k_4 b, k_3 b],$$

$$(H2) f(w) < 2k_2 b \text{ for } w \in [0, k_3 b],$$

$$(H3) f(w) < 8(k_3 b - k_5 b) \text{ for } w \in [0, k_3 b],$$

$$(H4) f(w) > 128 \left(k_4 b - \frac{192}{913} k_1 b \right) \text{ for } w \in [0, k_4 b], \text{ and}$$

$$(H5) f(w) < \frac{512}{55} \left(k_3 b - \frac{9}{256} k_2 b \right) \text{ for } w \in \left[\frac{1}{8} b (k_5 + 8k_1), k_3 b \right].$$

Then T has a fixed point $x^* \in A(\beta, k_3 b, \alpha, k_4 b)$; i.e. (1.1), (1.2) has a solution $x^* \in A(\beta, k_3 b, \alpha, k_4 b)$.

Proof. Let

$$A = \{x \in \mathcal{P} : x(3/16) - x(1/8) > k_1 b \text{ and } x'(0) < k_2 b\}.$$

Consider $A(\beta, k_3 b, \alpha, k_4 b) = \{x \in A : k_4 b < x(t) < k_3 b \text{ for } t \in [1/8, 1/2]\}$, and

$$A(\beta, k_3 b, \psi, k_5 b) = \{x \in A : x(t) < k_3 b \text{ for } t \in [0, 1/2] \text{ and } x'(1/4) > k_5 b\}.$$

(A1) We show $A(\beta, k_3b, \alpha, k_4b)$ is bounded. Let $x \in A(\beta, k_3b, \alpha, k_4b)$. Then

$$\begin{aligned} \|x\| &= \max_{t \in [0,1]} |x(t)| + \max_{t \in [0,1]} |x'(t)| \\ &= x(1/2) + x'(0) \leq k_3b + k_2b \\ &= (k_3 + k_2)b. \end{aligned}$$

Therefore, $A(\beta, k_3b, \alpha, k_4b)$ is bounded open subset of \mathcal{P} .

Next, we show $A(\beta, k_3b, \alpha, k_4b) \cap A(\beta, k_3b, \psi, k_5b) \neq \emptyset$. Choose K such that $K \leq 8k_3$, $K \geq \frac{128}{7}k_4$, and $K \geq \frac{8}{3}k_5$. Let

$$w_0(t) = \int_0^1 G(t, s) Kb \, ds.$$

Then $x \in \mathcal{P}$,

$$\begin{aligned} \beta(w_0) &= w_0\left(\frac{1}{2}\right) = \int_0^1 G\left(\frac{1}{2}, s\right) Kb \, ds \\ &= Kb \int_0^{1/2} s \, ds \\ &= \frac{Kb}{8} \\ &\leq k_3b. \end{aligned}$$

In addition,

$$\begin{aligned} \alpha(w_0) &= w_0\left(\frac{1}{8}\right) = \int_0^1 G\left(\frac{1}{8}, s\right) Kb \, ds \\ &= \int_0^{1/8} s \left(1 - \frac{1}{8}\right) Kb \, ds + \int_{1/8}^1 \frac{1}{8} (1 - s) Kb \, ds \\ &= Kb \left(\frac{7}{1024} + \frac{49}{1024} \right) \\ &= \frac{7Kb}{128} \\ &\geq k_4b. \end{aligned}$$

Lastly,

$$\begin{aligned}
\psi(w_0) &= w'_0 \left(\frac{1}{8} \right) = \int_0^{1/8} (-s)Kb \, ds + \int_{1/8}^1 (1-s)Kb \, ds \\
&= Kb \left(-\frac{1}{128} + \frac{49}{128} \right) \\
&= \frac{3Kb}{8} \\
&\geq k_5b.
\end{aligned}$$

Therefore,

$$w_0 \in A(\beta, k_3b, \alpha, k_4b) \cap A(\beta, k_3b, \psi, k_5b)$$

and

$$A(\beta, k_3b, \alpha, k_4b) \cap A(\beta, k_3b, \psi, k_5b) \neq \emptyset.$$

Finally, we show if $x \in \partial A \cap \overline{\mathcal{P}(\beta, k_3b, \alpha, k_4b)}$, then $Tx \neq x$. Let $x \in A(\beta, k_3b, \alpha, k_4b)$ with $Tx = x$. Then

$$x \left(\frac{1}{8} \right) > k_4b \text{ and } x \left(\frac{1}{2} \right) < k_3b.$$

Now,

$$\begin{aligned}
x \left(\frac{3}{16} \right) - x \left(\frac{1}{8} \right) &= (Tx) \left(\frac{3}{16} \right) - (Tx) \left(\frac{1}{8} \right) \\
&= \int_0^{3/16} s \left(1 - \frac{3}{16} \right) f(x(s)) \, ds + \int_{3/16}^1 \frac{3}{16} (1-s) f(x(s)) \, ds \\
&\quad - \int_0^{1/8} s \left(1 - \frac{1}{8} \right) f(x(s)) \, ds - \int_{1/8}^1 \frac{1}{8} (1-s) f(x(s)) \, ds \\
&= \int_0^{1/8} -\frac{1}{16} s f(x(s)) \, ds + \int_{1/8}^{3/16} \frac{1}{16} (15s - 2) f(x(s)) \, ds \\
&\quad + \int_{3/16}^1 \frac{11}{16} (1-s) f(x(s)) \, ds \\
&= \int_0^{1/8} -\frac{1}{16} s f(x(s)) \, ds + \int_{1/8}^{3/16} \frac{1}{16} (15s - 2) f(x(s)) \, ds \\
&\quad + \int_0^{13/16} \frac{11}{16} s f(x(s)) \, ds
\end{aligned}$$

$$\begin{aligned}
&= \int_0^{1/8} \frac{5}{8} s f(x(s)) ds + \int_{1/8}^{3/16} \frac{1}{8} (13s - 1) f(x(s)) ds \\
&\quad + \int_{3/16}^{13/16} \frac{11}{16} s f(x(s)) ds \\
&> \int_{1/8}^{3/16} \frac{1}{8} (13s - 1) f(x(s)) ds + \int_{3/16}^{13/16} \frac{11}{16} s f(x(s)) ds.
\end{aligned}$$

If $s \in [1/8, 13/16]$, $x(s) \in [k_4b, k_3b]$. By (H1), $f(w) > \frac{4,096}{913} k_1b$ for $w \in [k_4b, k_3b]$. Thus, $f(x(s)) > \frac{4,096}{913} k_1b$ for $s \in [1/8, 3/16]$. So,

$$\begin{aligned}
x\left(\frac{3}{16}\right) - x\left(\frac{1}{8}\right) &> \int_{1/8}^{3/16} \frac{1}{8} (13s - 1) \frac{4,096}{913} k_1b ds + \int_{3/16}^{13/16} \frac{11}{16} s \frac{4,096}{913} k_1b ds \\
&= \frac{4,096}{(8)(913)} k_1b \int_{1/8}^{3/16} (13s - 1) ds + \frac{(11)(4,096)}{(16)(913)} k_1b \int_{3/16}^{13/16} s ds \\
&= \frac{4,096}{7,304} k_1b \left(\frac{13s^2}{2} - s \right) \Big|_{1/8}^{3/16} + \frac{45,056}{14,608} k_1b \left(\frac{s^2}{2} \right) \Big|_{3/16}^{13/16} \\
&= \frac{512}{913} k_1b \left(\frac{13s^2}{2} - s \right) \Big|_{1/8}^{3/16} + \frac{256}{83} k_1b \left(\frac{s^2}{2} \right) \Big|_{3/16}^{13/16} \\
&= \frac{33}{913} k_1b + \frac{640}{664} k_1b \\
&= \frac{364k_1b + 7,040k_1b}{7,304} \\
&= \frac{7,304}{7,304} k_1b \\
&= k_1b.
\end{aligned}$$

Also, $x'(0) = (Tx)'(0) = \int_0^1 (1 - s) f(x(s)) ds$. If $s \in [0, 1]$, $x(s) \in [0, k_3]$. By (H2), $f(w) < 2k_2b$ for $w \in [0, k_3b]$. Thus, $f(x(s)) < 2k_2b$ for $s \in [0, 1]$. So,

$$\begin{aligned}
x'(0) &< \int_0^1 (1 - s) 2k_2b ds \\
&= 2k_2b \int_0^1 (1 - s) ds \\
&= 2k_2b \left(s - \frac{s^2}{2} \right) \Big|_0^1
\end{aligned}$$

$$\begin{aligned}
&= 2k_2b \left(1 - \frac{1}{2}\right) \\
&= 2k_2b \left(\frac{1}{2}\right) \\
&= k_2b.
\end{aligned}$$

Thus, $x \notin \partial A(\beta, k_3b, \alpha, k_4b)$. So by contrapositive, if $x \in \partial A(\beta, k_3b, \alpha, k_4b)$, then $Tx \neq x$.

(A2) Let $x \in \partial A(\beta, k_3b, \alpha, k_4b)$ with $\alpha(x) = k_4b$. Now $x\left(\frac{1}{8}\right) = k_4b$.

$$\begin{aligned}
\alpha(Tx) &= (Tx) \left(\frac{1}{8}\right) \\
&= \int_0^1 G\left(\frac{1}{8}, s\right) f(x(s)) ds \\
&= \int_0^{1/8} s \left(1 - \frac{1}{8}\right) f(x(s)) ds + \int_{1/8}^1 \frac{1}{8} (1-s) f(x(s)) ds \\
&= \int_0^{1/8} \frac{7}{8} s f(x(s)) ds + \int_{1/8}^1 \frac{1}{8} (1-s) f(x(s)) ds \\
&= \int_0^{1/8} \frac{7}{8} s f(x(s)) ds + \int_0^{7/8} \frac{1}{8} s f(x(s)) ds \\
&= \int_0^{1/8} s f(x(s)) ds + \int_{1/8}^{7/8} \frac{1}{8} s f(x(s)) ds.
\end{aligned}$$

If $s \in [0, 1/8]$, $x(s) \in [0, k_4b]$. By (H4), $f(w) > 128 \left(k_4b - \frac{192}{913}k_1b\right)$ for $w \in [0, k_4b]$. Thus, $f(x(s)) > 128 \left(k_4b - \frac{192}{913}k_1b\right)$ for $s \in [0, k_4b]$. Also, if $s \in [1/8, 7/8]$, $x(s) \in [k_4b, k_3b]$. By (H1), $f(w) > \frac{4,096}{913}k_1b$ for $w \in [k_4b, k_3b]$. Thus, $f(x(s)) > \frac{4,096}{913}k_1b$ for $s \in [1/8, 7/8]$. Therefore,

$$\begin{aligned}
\alpha(Tx) &> \int_0^{1/8} s \left(128 \left(k_4b - \frac{192}{913}k_1b\right)\right) ds + \int_{1/8}^{7/8} s \left(\frac{4,096}{913}k_1b\right) ds \\
&= 128 \left(k_4b - \frac{192}{913}k_1b\right) \int_0^{1/8} s ds + \frac{4,096}{913}k_1b \int_{1/8}^{7/8} \frac{1}{8} s ds \\
&= 128 \left(k_4b - \frac{192}{913}k_1b\right) \left(\frac{s^2}{2}\right) \Big|_0^{1/8} + \frac{4,096}{913}k_1b \left(\frac{s^2}{16}\right) \Big|_{1/8}^{7/8}
\end{aligned}$$

$$\begin{aligned}
&= 128 \left(k_4 b - \frac{192}{913} k_1 b \right) \left(\frac{1}{128} \right) + \frac{4,096}{913} k_1 b \left(\frac{1}{16} \left(\frac{49}{64} - \frac{1}{64} \right) \right) \\
&= k_4 b - \frac{192}{913} k_1 b + \frac{4,096}{913} k_1 b \left(\frac{3}{64} \right) \\
&= k_4 b - \frac{192}{913} k_1 b + \frac{192}{913} k_1 b \\
&= k_4 b.
\end{aligned}$$

(A3) Let $x \in \partial A(\beta, k_3 b, \alpha, k_4 b)$ with $\beta(x) = k_3 b$.

Suppose $\psi(x) \geq k_5 b$. Then $(x)' \left(\frac{1}{8} \right) \geq k_5 b$. Now $\psi(x) \geq k_5 b$ and x is increasing, so $x \left(\frac{1}{8} \right) \geq \frac{1}{8} k_5 b$, since

$$\begin{aligned}
\frac{1}{8} k_5 b &\leq \frac{\psi(x)}{8} \\
&= \int_0^{1/8} x' \left(\frac{1}{8} \right) ds \\
&\leq \int_0^{1/8} x'(s) ds \\
&= x \left(\frac{1}{8} \right).
\end{aligned}$$

Also, since $x \in A$,

$$\begin{aligned}
x \left(\frac{3}{16} \right) &\geq x \left(\frac{1}{8} \right) + k_1 b \\
&\geq \frac{1}{8} b (k_5 + 8k_1).
\end{aligned}$$

Thus,

$$\begin{aligned}
\beta(Tx) &= \int_0^1 G \left(\frac{1}{2}, s \right) f(x(s)) ds \\
&= \int_0^{1/2} s f(x(s)) ds \\
&= \int_0^{3/16} s f(x(s)) ds + \int_{3/16}^{1/2} s f(x(s)) ds.
\end{aligned}$$

If $s \in [0, 3/16]$, $x(s) \in [0, k_3 b]$. By (H2), $f(w) < 2k_2 b$ for $w \in [0, k_3 b]$. Thus, $f(x(s)) < 2k_2 b$ for $s \in [0, 3/16]$. Also, for $s \in [3/16, 1/2]$, $x(s) \in [\frac{1}{8} b (k_5 + 8k_1), k_3 b]$. By (H5), $f(w) < \frac{512}{55} (k_3 b - \frac{9}{256} k_2 b)$ for $w \in [\frac{1}{8} b (k_5 + 8k_1), k_3 b]$. Thus, $f(x(s)) <$

$\frac{512}{55} (k_3b - \frac{9}{256}k_2b)$ for $s \in [3/16, 1/2]$. Therefore,

$$\begin{aligned}
\beta(Tx) &< \int_0^{3/16} s \, 2k_2b \, ds + \int_{3/16}^{1/2} s \left(\frac{512}{55} \left(k_3b - \frac{9}{256}k_2b \right) \right) ds \\
&= 2k_2b \int_0^{3/16} s \, ds + \frac{512}{55} \left(k_3b - \frac{9}{256}k_2b \right) \int_{3/16}^{1/2} s \, ds \\
&= 2k_2b \left(\frac{9}{512} \right) + \frac{512}{55} \left(k_3b - \frac{9}{256}k_2b \right) \left(\frac{55}{512} \right) \\
&= \frac{9}{256}k_2b + k_3b - \frac{9}{256}k_2b \\
&= k_3b.
\end{aligned}$$

Next, suppose $\psi(Tx) < k_5b$. Now $(Tx)' \left(\frac{1}{8} \right) < k_5b$. So,

$$\begin{aligned}
\psi(Tx) &= (Tx)' \left(\frac{1}{8} \right) \\
&= \int_0^{1/8} (-s)f(x(s))ds + \int_{1/8}^1 (1-s)f(x(s))ds \\
&= \int_0^{1/8} (-s)f(x(s))ds + \int_0^{7/8} sf(x(s))ds \\
&= \int_{1/8}^{7/8} sf(x(s))ds \\
&< k_5b.
\end{aligned}$$

Thus,

$$\begin{aligned}
\beta(Tx) &= (Tx) \left(\frac{1}{2} \right) \\
&= \int_0^1 sf(x(s))ds \\
&= \int_0^{1/8} sf(x(s))ds + \int_{1/8}^{7/8} sf(x(s))ds + \int_{7/8}^1 sf(x(s))ds.
\end{aligned}$$

We also know that

$$\int_{7/8}^1 sf(x(s))ds = \int_0^{1/8} (1-s)f(x(s))ds.$$

So,

$$\int_0^{1/8} s f(x(s)) ds + \int_0^{1/8} (1-s) f(x(s)) ds = \int_0^{1/8} f(x(s)) ds.$$

Therefore, by combining them together, we get

$$\begin{aligned} \beta(Tx) &= \int_{1/8}^{7/8} s f(x(s)) ds + \int_0^{1/8} f(x(s)) ds \\ &< k_5 b + \int_0^{1/8} f(x(s)) ds \end{aligned}$$

If $s \in [0, 1/8]$, $x(s) \in [0, k_3 b]$. By (H3), $f(w) < 8(k_3 b - k_5 b)$ for $w \in [0, k_3 b]$. Thus, $f(x(s)) < 8(k_3 b - k_5 b)$ for $s \in [0, k_3 b]$. So,

$$\begin{aligned} \beta(Tx) &< k_5 b + \int_0^{1/8} 8(k_3 b - k_5 b) ds \\ &= k_5 b + 8(k_3 b - k_5 b) \Big|_0^{1/8} \\ &= k_5 b + 8(k_3 b - k_5 b) \left(\frac{1}{8}\right) \\ &= k_5 b + k_3 b - k_5 b \\ &= k_3 b. \end{aligned}$$

Therefore, T has a fixed point $x^* \in A(\beta, k_3 b, \alpha, k_4 b)$, and (1.1), (1.2) has a positive symmetric solution $x^* \in A(\beta, k_3 b, \alpha, k_4 b)$. \square

3.3 Example

Example 3.3. Example: Let $k_1 = 8380$, $k_2 = 230400$, $k_3 = 12140$, $k_4 = 1800$, and $k_5 = 80$. Then let us verify if the mentioned values satisfy the inequalities from (a) to (i).

(a) $\frac{128}{7}k_4 < 8k_3$. Then we have

$$\begin{aligned}\frac{128}{7}(1800) &< 8(12140) \\ 32914.2857 &< 97120.\end{aligned}$$

(b) $k_1 < \left(\frac{83}{40}\right) \left(k_3 - \frac{9}{256}k_2\right)$. Similarly,

$$8380 < \frac{83}{40} \left(12140 - \frac{9}{256}(230400)\right) = 8383.$$

(c) $16 \left(k_4 - \left(\frac{192}{913}\right)k_1\right) < k_3 - k_5$.

$$\begin{aligned}16 \left(1800 - \frac{192}{913}(8380)\right) &< 12140 - 80 \\ 16(1800 - 1762.278204) &< 12060 \\ 603.5487 &< 12060.\end{aligned}$$

(d) $\frac{4}{55} \left(k_3 - \left(\frac{9}{256}\right)k_2\right) > k_4 - \left(\frac{192}{913}\right)k_1$.

$$\begin{aligned}\frac{4}{55} \left(12140 - \frac{9}{256}(230400)\right) &> 1800 - \frac{192}{913}(8380) \\ \frac{4}{55}(4040) &> 1800 - 1762.278204 \\ 293 &> 37.7218\end{aligned}$$

(e) $k_2 > 64 \left(k_4 - \left(\frac{192}{913}\right)k_1\right)$

$$230400 > 64 \left(1800 - \frac{192}{913}(8380)\right) = 2414.1950$$

$$(f) \quad k_2 > \left(\frac{2048}{913}\right) k_1.$$

$$230400 > \frac{2048}{913}(8380) = 18797.6342$$

$$(g) \quad k_3 - k_5 > \left(\frac{512}{913}\right) k_1$$

$$12140 - 80 > \frac{512}{913}(8380)$$

$$12060 > 4699.4085$$

$$(h) \quad k_3 > \frac{1}{8}(k_5 + 8k_1).$$

$$12140 > \frac{1}{8}(80 + 8(8380)) = 8390.$$

$$(i) \quad k_4 < \left(\frac{1}{8}\right)(k_5 + 8k_1).$$

$$1800 < \frac{1}{8}(80 + 8(8380)) = 8390.$$

This shows that k_1 to k_5 satisfy (a) – (i). Thus, by letting $b = 1$ we have the following:

$$(H1) \quad f(w) > \frac{34324480}{913} \text{ for } w \in [1800, 12140],$$

$$(H2) \quad f(w) < 460800 \text{ for } w \in [0, 12140],$$

$$(H3) \quad f(w) < 96480 \text{ for } w \in [0, 12140],$$

$$(H4) \quad f(w) > \frac{4408320}{913} \text{ for } w \in [0, 1800], \text{ and}$$

$$(H5) \quad f(w) < \frac{413696}{11} \text{ for } w \in [8390, 12140].$$

Then the constant function $f(w) = 37600$ satisfies the conditions mentioned above.

Therefore, Theorem 3.1 guarantees the existence x^* of a positive symmetric solution

of the boundary value problem

$$x'' + 37600 = 0,$$

$$x(0) = 0 = x(1),$$

with $x^* \in A(\beta, 12140, \alpha, 1800)$.

BIBLIOGRAPHY

- [1] Abdulmalik Al Twaty and Paul W. Eloe. Concavity of solutions of a $2n$ -th order problem with symmetry. *Opuscula Math.*, 33(4):603–613, 2013.
- [2] Abdulmalik A. Altwaty and Paul Eloe. The role of concavity in applications of Avery type fixed point theorems to higher order differential equations. *J. Math. Inequal.*, 6(1):79–90, 2012.
- [3] Douglas Anderson, Richard Avery, Johnny Henderson, and Xueyan Liu. Multiple fixed point theorems utilizing operators and functionals. *Commun. Appl. Anal.*, 16(3):377–387, 2012.
- [4] Douglas R. Anderson, Richard I. Avery, and Johnny Henderson. Functional expansion—compression fixed point theorem of Leggett-Williams type. *Electron. J. Differential Equations*, pages No. 63, 9, 2010.
- [5] Douglas R. Anderson, Richard I. Avery, Johnny Henderson, and Xueyan Liu. Operator type expansion-compression fixed point theorem. *Electron. J. Differential Equations*, pages No. 42, 11, 2011.
- [6] Douglas R. Anderson, Richard I. Avery, Johnny Henderson, and Xueyan Liu. Fixed point theorem utilizing operators and functionals. *Electron. J. Qual. Theory Differ. Equ.*, pages No. 12, 16, 2012.
- [7] Richard Avery, Douglas Anderson, and Johnny Henderson. Some fixed point theorems of Leggett-Williams type. *Rocky Mountain J. Math.*, 41(2):371–386, 2011.
- [8] Richard Avery, Paul Eloe, and Johnny Henderson. A Leggett-Williams type theorem applied to a fourth order problem. *Commun. Appl. Anal.*, 16(4):579–588, 2012.

- [9] Richard Avery, Johnny Henderson, and Donal O'Regan. Functional compression-expansion fixed point theorem. *Electron. J. Differential Equations*, pages No. 22, 12, 2008.
- [10] Richard I. Avery, Douglas R. Anderson, and Johnny Henderson. An extension of the compression-expansion fixed point theorem of functional type. *Electron. J. Differential Equations*, pages Paper No. 253, 12, 2016.
- [11] Richard I. Avery, Douglas R. Anderson, and Johnny Henderson. Generalization of the functional omitted ray fixed point theorem. *Comm. Appl. Nonlinear Anal.*, 25(1):39–51, 2018.
- [12] Richard I. Avery, John R. Graef, and Xueyan Liu. Compression fixed point theorems of operator type. *J. Fixed Point Theory Appl.*, 17(1):83–97, 2015.
- [13] Richard I. Avery, Johnny Henderson, and Douglas R. Anderson. Existence of a positive solution to a right focal boundary value problem. *Electron. J. Qual. Theory Differ. Equ.*, pages No. 5, 6, 2010.
- [14] Richard W. Leggett and Lynn R. Williams. Multiple positive fixed points of nonlinear operators on ordered Banach spaces. *Indiana Univ. Math. J.*, 28(4):673–688, 1979.